



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Experimental evidence that clay inhibits bacterial decomposers: Implications for preservation of organic fossils

Citation for published version:

McMahon, S, Anderson, RP, Saupe, EE & Briggs, DEG 2016, 'Experimental evidence that clay inhibits bacterial decomposers: Implications for preservation of organic fossils', *Geology*, vol. 44, no. 10, pp. 867-870. <https://doi.org/10.1130/G38454.1>

Digital Object Identifier (DOI):

[10.1130/G38454.1](https://doi.org/10.1130/G38454.1)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Geology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 Experimental evidence that clay inhibits bacterial decomposers,
2 with implications for the preservation of organic fossils

3

4 **Sean McMahon, Ross P. Anderson, Erin E. Saupe, and Derek E. G. Briggs**

5 Department of Geology and Geophysics, Yale University, 210 Whitney Avenue, New
6 Haven, CT 06511, USA

7

8 **ABSTRACT**

9

10 Exceptionally preserved organic fossils are commonly associated with clay-rich horizons
11 or directly with clay minerals. It has been posited that interactions between clay minerals
12 and organic tissues inhibit enzymatic reactions or protect carcasses in such a way that
13 decay is inhibited. However, interactions between clay minerals and the biological agents
14 of decay, especially bacteria, may be at least as important to preservation potential. Here
15 we show that clays of particle size $< 2 \mu\text{m}$ in suspensions exceeding 10 mg/ml in
16 concentration inhibit the growth of *Pseudoalteromonas luteoviolacea*, a marine
17 heterotrophic bacterium involved in the decay of marine animals. Such clay–microbe
18 interactions can contribute to exceptional preservation and specific examples may play a
19 role in shaping the distribution of Konservat-Lagerstätten through time.

20

INTRODUCTION

Exceptionally preserved fossil assemblages (Konservat-Lagerstätten *sensu* Seilacher, 1970) that yield the remains of soft tissues constitute an invaluable source of information about the history of life on Earth. In diverse lagerstätten of all ages, organic remains are associated with aluminosilicate clay minerals (e.g., Butterfield, 1994; Anderson et al. 2011; Laflamme et al., 2011; Cai et al., 2012; Pan et al., 2014; Wacey et al., 2014). A large and paleobiologically important subset of these lagerstätten exhibit Burgess-Shale type (BST) preservation, in which organic material is retained and flattened parallel to bedding in fine-grained siliciclastic lithologies often with high clay-to-organic ratios (Butterfield, 1990; Butterfield et al., 1994, Butterfield, 1995; Gaines, 2014). Indeed this is the principal mode of preservation for eukaryotic remains in Proterozoic successions (e.g. Cohen and Macdonald, 2015; Yuan et al., 2011; Butterfield et al., 1994), as well as soft-bodied Cambrian marine metazoan faunas (e.g., Butterfield, 1990; Gaines et al., 2008; Gaines 2014). Many workers have argued that temporal and environmental distributions of clay minerals impart patterns and biases to the fossil record (e.g. Butterfield, 1995). Butterfield (1990) attributed the preservation of organic material to the effect of clays in adsorbing and deactivating autolytic enzymes, Orr et al. (1998) proposed that early diagenetic clays templated original organic tissues, and Petrovich (2001) suggested that Fe^{2+} ions stabilized the tissues and acted as nucleation sites for clay minerals. Wilson and Butterfield (2014) proposed that Al^{3+} ions derived from a clay-rich substrate protected the tissues from decay in a process analogous to tanning and speculated that Fe-rich clays could have a similar effect. Independent of the association

of clays and fossils, there is abundant evidence that clay minerals interfere with bacterial growth (e.g. Wong et al., 2004; Williams et al., 2011; Morrison et al., 2016).

We tested the hypothesis that clay mineral particles inhibit the growth of bacterial decomposers by comparing the growth of *Pseudoalteromonas luteoviolacea* (a heterotrophic marine γ -proteobacterium) in the presence of various clay minerals, calcite, and in a mineral-free control. Bacteria of this widespread genus are commonly found in biofilms associated with live marine animals, have the capacity to degrade a range of tissue types, and constitute a large proportion of the bacterial assemblage in submerged carcasses (Skovhus et al., 2007; Raff et al., 2008; Dickson et al., 2011). The strain used here was shown to decompose sea urchin embryos *in vitro* (Raff et al., 2013).

MATERIALS AND METHODS

Five minerals were used in the experiments: berthierine (Scarborough, UK; Yale Peabody Museum (YPM) MIN 100586), calcite (Budapest, Hungary; YPM MIN 056322), glauconite (Odessa, Delaware; YPM MIN 043086), illite (Silver Hill, Montana; Clay Minerals Society #IMt-2), and kaolinite (Santa Cruz Biotechnology Company, USA). Specimens were rinsed in distilled water, ground with an agate mill and sonicated to obtain fine powders, which were then rinsed again by centrifuging in distilled water at $2773 \times g$ (4000 rpm) for 15 minutes, three times. Following 10 minutes of sonication to disaggregate particles, grains of up to two microns in diameter were obtained by centrifuging in distilled water at $97 \times g$ (750 rpm) for 5 minutes, discarding the pellet and retaining the supernatant, which was removed and dried at 45 °C. This ensured that all minerals were similar in particle size to natural mud and to each other.

Bacterial growth medium (Difco Marine Broth [MB]) was prepared in sterile conditions, filtered to remove particles >20 µm, and added in aliquots of 950 µl to glass culture vials (8 ml; Wheaton/Fisher Scientific) to which the clays were added immediately prior to autoclaving. Additional sonication was employed to disaggregate mineral particles prior to inoculation. Vials were prepared with suspended mineral concentrations of 5, 10 and 25 mg/ml (Fig. 1) for each of the five mineral species and for a control with no suspended mineral particles.

The inoculum was taken from a stock culture on agar plates of *Pseudoalteromonas luteoviolacea* (ATCC33492) and grown in MB to an optical density (OD₆₀₀) of ~0.5 (above sterile MB). Vortex-mixed aliquots of 50 µl were transferred under sterile conditions to each of the clay-bearing vials, which were incubated on a rotary shaker at 25 °C. The use of continuously agitated suspensions rather than settled sediment ensured thorough and homogeneous mixing of mineral particles and bacteria. Preliminary results suggested that the presence of mineral particles interfered with normal techniques for measuring bacterial growth. We therefore subsampled the experimental cultures (post vortexing) after 24 hours' growth and diluted them in fresh medium so that both clay particles and bacteria were 100 times less concentrated. The bacteria multiplied in the fresh medium so that geometric differences in population size were inherited from the original cultures, while clay concentration remained minimal. Turbidity at 600 nm (a standard proxy for bacterial cell density) was measured using a spectrophotometer (Hach DR/2010) in mid-exponential phase (after six hours of growth), at which time the disparity in population sizes was clearly evident. Initial values were subtracted so that any turbidity due to clay was excluded. All experiments were performed in quadruplicate.

Statistical analyses were performed using the R programming language (R Core Team, 2016). For each mineral concentration turbidity was normalized by the mean of the clay-free control treatment. Using the control-standardized data, two-way analyses of variance (ANOVA, Type III Sum of Squares) were performed on turbidity levels in the ‘car’ R package (Fox and Weisberg, 2011), with mineral concentration and mineral species as factors. Planned post-hoc comparisons were performed in the phia R package (De Rosario-Martinez, 2013) using the Holm (1979) correction for multiple comparisons. Such pair-wise comparisons were made to test for statistical differences between factor levels (e.g., kaolinite versus control) and were made independently both within and across the three mineral concentrations. Effect sizes were calculated using the etaSquared function in the lsr R package (Navarro, 2015), which reveals the proportion of the variance in the dependent variable that is attributable to the factor in question. We also examined the effect of mineral species and mineral concentration on bacterial growth by considering the four clay minerals (berthierine, glauconite, illite, and kaolinite) as a single category. The above analyses were repeated using this binned dataset. All data were normally distributed, homoscedastic, and showed linear structure based on Q-Q, scale-location, and residuals vs. fitted values plots, respectively.

RESULTS

A two-factor analysis of variance showed significant main effects of mineral concentration, $F(2, 53) = 58.3$, $p < 0.05$ ($\eta^2 = 0.40$) and mineral species, $F(5, 53) = 2.9$, $p < 0.05$ ($\eta^2 = 0.29$) on turbidity, but these effects were qualified by an interaction between

the two variables, $F(10, 53) = 11.56$, $p < 0.05$ ($\eta^2 = 0.20$) (Supplementary Table 1), discussed below.

Post-hoc comparison tests revealed no significant difference in bacterial growth among mineral species compared to the control at 5 mg/ml, with the exception of glauconite (Fig. 1A; Supplementary Table 2). The significance of the response to glauconite at 5 mg/ml suggests an unusually low minimum inhibitory concentration with just a slightly increased effect at higher concentrations. Mineral suspensions of 10 and 25 mg/ml resulted in significantly depressed bacterial growth for all mineral species compared to the clay-free control, with the exception of calcite (Figs. 1A and 1B). The response at 25 mg/ml is much more pronounced to berthierine and kaolinite than to the other clay minerals—a fourfold decrease in turbidity compared to the control and threefold compared to calcite. We observed an inverse relationship between mineral concentration and turbidity for all mineral species except calcite, but the strength and significance of this relationship varies with mineral species (Supplementary Table 3). Although calcite appears to impede bacterial growth more strongly at 10 mg/ml than at 25 mg/ml, this is not significant.

In order to examine the impact of clays in general, we compared the effect of the clay minerals as a group with that of calcite and the control (Fig. 2). The two-factor analysis of variance showed significant main effects of mineral concentration, $F(2, 62) = 5.98$, $p < 0.05$ ($\eta^2 = 0.40$) and mineral species, $F(2, 62) = 1.23$, $p < 0.05$ ($\eta^2 = 0.30$) on turbidity, qualified by their interaction, $F(4, 62) = 12.246$, $p < 0.05$ ($\eta^2 = 0.13$) (Supplementary Table 1). Post hoc comparisons revealed no significant effect of clay or calcite on bacterial growth at 5 mg/ml. Both clay and calcite minerals, however, affected growth significantly at 10 and 25 mg/ml, but in both instances clay minerals were

associated with significantly less growth than calcite (Supplementary Table 2). Increasing clay concentration resulted in significantly depressed bacterial growth, but the same was not true for increasing calcite concentration (Supplementary Table 3). The marginal means and standard deviations of all within-concentration analyses are presented in Supplementary Table 4.

In summary, the results of our experiments show that the growth of *Pseudoalteromonas luteoviolacea*, a marine heterotrophic bacterium known to degrade animal tissues, is significantly inhibited by the presence of clay-sized mineral particles. With increasing mineral concentration, this inhibition becomes more pronounced and shows greater variation between mineral species. Clay minerals produce a stronger effect than calcite particles of similar size.

DISCUSSION AND CONCLUSIONS

Our experimental results support the hypothesis that clay minerals impede the growth of bacterial decomposers, providing a way to facilitate the preservation of soft tissues. Hypotheses to explain particular instances of exceptional fossil preservation have implicated both detrital clays, and authigenic clays that formed in sedimentary pore spaces and in association with decaying organic matter (e.g., Butterfield, 1990; Orr et al. 1998; Gabbott et al. 2001). Our results are relevant in both cases, and furthermore they suggest that certain clays are more likely to inhibit bacterial activity than others.

Our results are consistent with the usually high clay content of Cambrian Burgess Shale-type fossil-hosting sediments (Curtin and Gaines, 2011; Forchielli et al., 2014; Gaines et al., 2011; Powell, 2003). However, there is currently limited documentation of

associations with specific clay compositions. An exception is the Mount Cap Formation, where glauconite is associated with fossiliferous mudrock horizons (Aitken et al., 1973; Butterfield, 1994). Indeed, macrostratigraphic data from Laurentia reveal a spike in glauconite production during the Cambrian (Peters and Gaines, 2012). More generally, it has been argued that the rise and fall of BST preservation tracked the weathering flux of Al-rich clays as well as particular biogeochemical conditions that affected the composition of clay in marine sediment (Butterfield, 1995; Wilson & Butterfield 2014). Preliminary data through a Cryogenian succession in Mongolia provide a further example of clays of specific composition (in this case berthierine) in fossiliferous horizons (Anderson et al., 2014).

In both BST and non-BST lagerstätten, clay minerals can also be found intimately associated with organic fossils, often coating their external surfaces (e.g. Gabbott, 1998; Orr et al., 1998; Gabbott et al., 2001; Anderson et al., 2011; Laflamme et al., 2011; Cai et al., 2012; Pan et al., 2014; Wacey et al., 2014). A more refined mineralogical characterisation of clays within fossiliferous laminae and surrounding individual fossils awaits the application of emerging microscopic techniques (Tosca et al., 2015).

Our results are also consistent with the higher quality of organic tissue preservation associated with clay minerals as opposed to carbonate throughout the geological record. Fine-grained carbonates tend to preserve organic tissues more rarely and with less fidelity than clays (e.g., Butterfield et al., 1994). Organic fossils in Proterozoic carbonates, for example, tend to be dominated by robust testate forms (e.g., Bosak et al., 2011a; Bosak et al., 2011b; Bosak et al., 2012; Dalton et al., 2013) in contrast to more delicate forms preserved in clay-rich shales (e.g. Butterfield et al., 1994; Butterfield & Rainbird, 1998).

However, other factors may contribute to these differences, such as the higher sedimentation rate represented by clastic beds (Canfield, 1994).

The antibacterial properties of clays are attributed mostly to the toxicity of metal cations, particularly Al^{3+} and Fe^{2+} (e.g., Wong et al., 2004; Morrison et al., 2016). Diverse bacteria are susceptible to Al^{3+} , while excessive Fe^{2+} in aerobic conditions causes oxidative damage to bacterial cells (Guida et al., 1991; Kapoor and Arora, 1998; Amonette et al., 2003; Imlay et al., 2008). It is therefore striking that kaolinite and berthierine, the most aluminum-rich and ferrous-iron-rich clays in our experiments respectively, were associated with the strongest suppression of bacterial growth at 25 mg/ml. Kaolinite, which has been shown to preserve experimentally buried invertebrate carcasses better than quartz, calcite, and montmorillonite, also inhibits the growth of sulfate-reducing bacteria, autotrophic methanogens, and a heterotrophic soil bacterium (Wong et al., 2004; Wu et al., 2013; Wilson and Butterfield, 2014; Liu et al., 2016). Natural ‘blue’ clays, which release both Al^{3+} and Fe^{2+} from illite-smectite into solution, are effective antibiotic agents (Morrison et al., 2016); these two ions work synergistically to disrupt and oxidatively damage bacterial cells.

Our results show that clays impede the growth of decay bacteria, providing clear evidence that they are likely to have played a role in promoting organic preservation (Butterfield, 1990; Petrovich, 2001; Wilson and Butterfield, 2014). However, clay–bacterial interactions are highly specific with respect to both clay mineral composition and bacterial strain. Na-montmorillonite, for example, inhibits sulfate reducers but increases the longevity, growth rate and metabolic activity of several other groups of decomposers, probably due to its tendency to adsorb trace metals from the environment (Kunc & Stotzky, 1974; Hwang and Tate, 1997; Wong et al., 2004; Wu et al., 2013). The

chemical changes associated with decomposition are likely to affect the leaching and adsorption behavior of the clay as well as the speciation and toxicity of leached ions (Guida et al., 1991; Andrews et al., 2003; Morrison et al., 2014). Further experiments involving a wider range of relevant strains, mineral species, and environmental conditions are required to unravel the role of clay–microbe interactions in exceptional preservation.

ACKNOWLEDGMENTS

Financial support was provided by the NASA Astrobiology Institute NNA13AA90A, *Foundations of Complex Life, Evolution, Preservation and Detection on Earth and Beyond*. Stefan Nicolescu (Yale Peabody Museum) provided mineral specimens and documentation, and R. A. Raff and M. E. Andrews provided bacterial strains and advice about their cultivation. Adrien Gau assisted with maintaining bacterial cultures. Nicholas Tosca provided valuable discussion. RPA was supported by NASA Headquarters under the Earth and Space Science Fellowship program (NNX14AP10H).

REFERENCES

Aitken, J.D., Macqueen, R.W., and Usher, J.L., 1973, Reconnaissance studies of Proterozoic and Cambrian stratigraphy, lower Mackenzie River area, District of Mackenzie: Geological Survey of Canada Papers, v. 73, no. 9, p. 1–163.

233 Amonette, J.E., Russell, C.K., Carosino, K.A., Robinson, N.L., and Ho, J.T., 2003,
 234 Toxicity of Al to *Desulfovibrio desulfuricans*: Applied and environmental microbiology,
 235 v. 69, no. 7, p. 4057–4066, doi:10.1128/AEM.69.7.4057-4066.2003.
 236
 237 Anderson, E.P., Schiffbauer, J.D., and Xiao, S.H., 2011, Taphonomic study of Ediacaran
 238 organic-walled fossils confirms the importance of clay minerals and pyrite in Burgess
 239 Shale-type preservation: *Geology*, v. 39, no. 7, p.643–646, doi:10.1130/G31969.1
 240
 241 Anderson, R.P., Macdonald, F.A., Tosca, N.J., Bosak, T., Bold, U., and Briggs, D.E.G.,
 242 2014, Taphonomy of eukaryotic microfossils between Cryogenian ice ages explored in
 243 the Zavkhan terrane, southwestern Mongolia: *Geological Society of America Abstracts*
 244 *with Programs*, v. 46, no. 6, p. 542.
 245
 246 Andrews, S.C., Robinson, A.K., and Rodríguez-Quñones, F., 2003, Bacterial iron
 247 homeostasis: *FEMS microbiology reviews*, v. 27, no. 2–3, p. 215–237,
 248 doi:10.1016/S0168-6445(03)00055-X.
 249
 250 Bosak, T., Lahr, D.J.G., Pruss, S.B., Macdonald, F.A., Dalton, L., and Matys, E.D.,
 251 2011a, Agglutinated tests in post-Sturtian cap carbonates of Namibia and Mongolia:
 252 *Earth and Planetary Science Letters*, v. 308, no. 1–2, p. 29–40,
 253 doi:10.1016/j.epsl.2011.05.030.
 254

255 Bosak, T., Lahr, D.J.G., Pruss, S.B., Macdonald, F.A., Gooday, A.J., Dalton, L., and
 256 Matys, E.D., 2012, Possible early foraminiferans in post-Sturtian (716-635 Ma) cap
 257 carbonates: *Geology*, v. 40, no. 1, p. 67–70, doi:10.1130/G32535.1.
 258
 259 Bosak, T., Macdonald, F.A., Lahr, D.J.G., and Matys, E.D., 2011b, Putative Cryogenian
 260 ciliates from Mongolia: *Geology*, v. 39, no. 12, p. 1123–1126, doi:10.1130/G32384.1.
 261
 262 Butterfield, N.J., 1990, Organic preservation of non-mineralizing organisms and the
 263 taphonomy of the Burgess Shale: *Paleobiology*, v. 16, no. 3, p. 272–286,
 264 doi:10.1017/S0094837300009994.
 265
 266 Butterfield, N.J., 1994, Burgess Shale-type fossils from a Lower Cambrian shallow-shelf
 267 sequence in northwestern Canada: *Nature*, v. 369, p. 477–479, doi:10.1038/369477a0.
 268
 269 Butterfield, N.J., 1995, Secular distribution of Burgess-Shale-type preservation: *Lethaia*,
 270 v. 28, no. 1, p. 1–13, doi: 10.1111/j.1502-3931.1995.tb01587.x.
 271
 272 Butterfield, N.J., and Harvey, T.H.P., 2012, Small carbonaceous fossils (SCFs): A new
 273 measure of early Paleozoic paleobiology: *Geology*, v. 40, no. 1, p. 71–74,
 274 doi:10.1130/G32580.1.
 275
 276 Butterfield, N. J., Knoll, A. H., and Swett, K., 1994, Paleobiology of the Neoproterozoic
 277 Svanbergfjellet Formation, Spitsbergen: *Fossils and Strata*, v. 34, p. 1–84.
 278

279 Butterfield, N. J., and Rainbird, R. H., 1998, Diverse organic-walled fossils, including
 280 “possible dinoflagellates” from the early Neoproterozoic of arctic Canada. *Geology*, v.
 281 26, p. 963–966, doi:10.1130/0091-7613(1998).
 282

283 Cai, Y., Schiffbauer, J.D., Hua, H., and Xiao, S.H., 2012, Preservational modes in the
 284 Ediacaran Gaijiashan lagerstätte: Pyritization, aluminosilification, and carbonaceous
 285 compression: *Paleogeography, Palaeoclimatology, Palaeoecology*, v. 326-328, p.109–
 286 117, doi:10.1016/j.palaeo.2012.02.009.
 287

288 Cohen, P.A., and Macdonald, F.A., 2015, The Proterozoic record of eukaryotes:
 289 *Paleobiology*, p. 1–23, doi:10.1017/pab.2015.25.
 290

291 Curtin, L.G., and Gaines, R.R., 2011, Burgess Shale-type preservation and detrital clay
 292 mineralogy: A test of the "reactive clay" hypothesis: *Geological Society of America*
 293 *Abstracts with Programs*, v. 43, no. 5, p. 108.
 294

295 Dalton, L.A., Bosak, T., Macdonald, F.A., Lahr, D.J.G., and Pruss, S.B., 2013,
 296 Preservational and morphological variability of assemblages of agglutinated eukaryotes
 297 in Cryogenian cap carbonates of northern Namibia: *Palaios*, v. 28, no. 1–2, p. 67–79,
 298 doi:10.2110/palo.2012.p12-084r.
 299

300 De Rosario-Martinez, H., 2013, phia: Post-Hoc Interaction Analysis. R Package Version
 301 0.2-1. <http://CRAN.R-project.org/package=phia>.
 302

303 Dickson, G.C., Poulter, R.T., Maas, E.W., Probert, P.K., and Kieser, J.A., 2011, Marine
 304 bacterial succession as a potential indicator of postmortem submersion interval: Forensic
 305 Science International, v. 209, no. 1, p.1–10, doi:10.1016/j.forsciint.2010.10.016.
 306
 307 Forchielli, A., Steiner, M., Kasbohm, J., Hu, S., and Keupp, H., 2014, Taphonomic traits
 308 of clay-hosted early Cambrian Burgess Shale-type fossil lagerstätten in South China:
 309 Palaeogeography, Palaeoclimatology, Palaeoecology, v. 398, p. 59–85,
 310 doi:10.1016/j.palaeo.2013.08.001.
 311
 312 Fox, J., and Weisberg, S., (2011). An {R} Companion to Applied Regression, Second
 313 Edition. Thousand Oaks CA: Sage. URL:
 314 <http://socserv.socsci.mcmaster.ca/jfox/Books/Companion>
 315
 316 Gabbott, S.E, 1998, Taphonomy of the Ordovician Soom Shale lagerstätte: An example
 317 of soft tissue preservation in clay minerals: Palaeontology, v. 41, p. 631–667.
 318
 319 Gabbott, S.E., Norry, M.J., Aldridge, R.J., Theron, J.N., 2001, Preservation of fossils in
 320 clay minerals; a unique example from the Upper Ordovician Soom Shale, South Africa:
 321 Proceedings of the Yorkshire Geological Society, v. 53, p. 237–244.
 322
 323 Gaines, R.R., Briggs, D.E.G. and Zhao, Y.-L. 2008 Cambrian Burgess Shale-type
 324 deposits share a common mode of fossilization. Geology, v. **36**, p. 755–758, doi:
 325 10.1130/G24961A.1.

326 Gaines, R.R., 2014, Burgess Shale-type preservation and its distribution in space and
 327 time, in Reading and Writing of the Fossil Record: Preservational Pathways to
 328 Exceptional Fossilization, Paleontological Society Papers, v. 20, p. 123–146.

329 Gaines, R.R., Mering, J.A., Zhao, Y., and Peng, J., 2011, Stratigraphic and microfacies
 330 analysis of the Kaili Formation, a candidate GSSP for the Cambrian Series 2-Series 3
 331 boundary: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 311, p. 171–183,
 332 doi:10.1016/j.palaeo.2011.08.018.

333 Guida, L., Saidi, Z., Hughes, M.N., and Poole, R. K., 1991, Aluminium toxicity and
 334 binding to *Escherichia coli*: Archives of Microbiology, v. 156, no. 6, 507–512,
 335 doi:10.1007/BF00245400.

336 Holm, S, 1979, A simple sequentially rejective multiple test procedure: Scandinavian
 337 Journal of Statistics, v. 6, p. 65–70.

338 Hwang, S., and Tate III, R. L., 1997, Interactions of clay minerals with *Arthrobacter*
 339 *crystallopoietes*: starvation, survival and 2-hydroxypyridine catabolism: Biology and
 340 Fertility of Soils, v. 24, p. 335–340, doi:10.1007/s003740050253.

341 Imlay, J. A., Chin, S. M., and Linn, S., 2008, Toxic DNA damage by hydrogen peroxide
 342 through the Fenton reaction in vivo and in vitro: Science, v. 240, p. 640–642,
 343 doi:10.1126/science.2834821.

344 Kapoor, K., and Arora, L., 1998, Aluminium induced toxicity and growth responses of
 345 cyanobacteria: Pollution Research, v. 17, no. 1, p. 25–31

346 Kunc, F., and Stotzky, G., 1974, Effect of clay minerals on heterotrophic microbial
 347 activity in soil: Soil Science, v. 118, no. 3, p. 186–195.

348 Laflamme, M., Schiffbauer, J.D., Narbonne, G.M., and Briggs, D.E.G., 2011, Microbial
 349 biofilms and the preservation of the Ediacara biota: *Lethaia*, v. 44, no. 2, p. 203–213, doi:
 350 10.1111/j.1502-3931.2010.00235.x.

351 Liu, D., Dong, H., Agrawal, A., Singh, R., Zhang, J., and Wang, H., Inhibitory effect of
 352 clay mineral on methanogenesis by *Methanosarcina mazei* and *Methanothermobacter*
 353 *thermautotrophicus*: *Applied Clay Science*, v. 126, p. 25–32,
 354 doi:10.1016/j.clay.2016.02.030 0169-1317/.

355 Morrison, K.D., Underwood, J.C., Metge, D.W., Eberl, D.D., and Williams, L.B., 2014.
 356 Mineralogical variables that control the antibacterial effectiveness of a natural clay
 357 deposit: *Environmental Geochemistry and Health*, v. 36, n. 4, p. 613–631,
 358 doi:10.1007/s10653-013-9585-0.

359 Morrison, K.D., Misra, R., and Williams, L.B., 2016, Unearthing the antibacterial
 360 mechanism of medicinal clay: A geochemical approach to combating antibiotic
 361 resistance: *Scientific Reports*, v. 6, doi:10.1038/srep19043.

362 Navarro, D. J., 2015, Learning statistics with R: A tutorial for psychology students and
 363 other beginners. (Version 0.5) University of Adelaide. Adelaide, Australia.

364 Orr, P.J., Briggs, D.E.G., and Kearns, S.L., 1998, Cambrian Burgess Shale animals
 365 replicated in clay minerals: *Science*, v. 281, no. 5380, p. 1173–1175,
 366 doi:10.1126/science.281.5380.1173.

367 Pan, Y., Sha, J., and Fuerisch, F.T., 2014, A model for organic fossilization of the Early
 368 Cretaceous Jehol Lagerstätte based on the taphonomy of “*Ephemeropsis trisetalis*”:
 369 *Palaos*, v. 29, no. 7, p. 363–377, doi:10.2110/palo.2013.119.

370 Peters, S.E., and Gaines, R.R., 2012, Formation of the Great Unconformity as a trigger
 371 for the Cambrian explosion: *Nature*, v. 484, no. 7394, p. 363–366,
 372 doi:10.1038/nature10969.

373 Petrovich, R., 2001, Mechanisms of fossilization of the soft-bodied and lightly armored
 374 faunas of the Burgess Shale and of some other classical localities: *American Journal of*
 375 *Science*, v. 301, no. 8, p. 683–726.

376 Powell, W., 2003, Greenschist-facies metamorphism of the Burgess Shale and its
 377 implications for models of fossil formation and preservation: *Canadian Journal of Earth*
 378 *Sciences*, v. 40, p. 13-25, doi:10.1139/e02-103.

379 R Core Team (2016). R: A language and environment for statistical computing. R
 380 Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

381 Raff, E.C., Schollaert, K.L., Nelson, D.E., Donoghue, P.C., Thomas, C.W., Turner, F.R.,
 382 Stein, B.D., Dong, X., Bengtson, S., Huldtgren, T., and Stampanoni, M., 2008, Embryo
 383 fossilization is a biological process mediated by microbial biofilms: *Proceedings of the*
 384 *National Academy of Sciences*, v. 105, no. 49, p. 19360–19365, doi:
 385 10.1073/pnas.0810106105.

386 Raff, E.C., Andrews, M.E., Turner, F.R., Toh, E., Nelson, D.E., and Raff, R.A., 2013,
 387 Contingent interactions among biofilm- forming bacteria determine preservation or
 388 decay in the first steps toward fossilization of marine embryos: *Evolution and*
 389 *Development*, v. 15, no. 4, p. 243–256, DOI: 10.1111/ede.12028.

390 Seilacher, A., 1970. Begriff und Bedeutung der Fossil-Lagerstätten. *Neues Jahrbuch für*
 391 *Geologie und Paläontologie, Monatshefte*, 1970, p. 34–39.

392 Skovhus, T.L., Holmström, C., Kjelleberg, S., and Dahllöf, I., 2007, Molecular
 393 investigation of the distribution, abundance and diversity of the genus
 394 *Pseudoalteromonas* in marine samples: FEMS microbiology ecology, v. 61, no. 2, p.
 395 348–361, doi: 10.1111/j.1574-6941.2007.00339.x.

396 Tosca, N.J., Pruss, S.B., and Strauss, J.V., 2015, A sedimentary fingerprint of oscillating
 397 redox through the Cambrian Period: Geological Society of America Abstracts with
 398 Programs, v. 47, no. 7, p. 269.

399 Wacey, D., Saunders, M., Roberts, M., Menon, S., Green, L., Kong, C., Culwick, T.,
 400 Strother, P., and Brasier, M.D., 2014, Enhanced cellular preservation by clay minerals in
 401 1 billion-year-old lakes: Scientific Reports, v. 4, 5841, doi:10.1038/srep05841.

402 Williams, L.B., Metge, D.W., Eberl, D.D., Harvey, R.W., Turner, A.G., Prapaipong, P.,
 403 and Poret-Peterson, A.T., 2011, What makes a natural clay antibacterial?: Environmental
 404 Science and Technology, v. 45, no. 8, p. 3768–3773, doi: 10.1021/es1040688.

405 Wilson, L.A., and Butterfield, N.J., 2014, Sediment effects on the preservation of
 406 Burgess Shale-type compression fossils: Palaios, v. 29, no. 4, p. 145–154, doi:
 407 10.2110/palo.2013.075.

408 Wong, D., Suflita, J.M., McKinley, J.P., and Krumholz, L.R., 2004, Impact of clay
 409 minerals on sulfate-reducing activity in aquifers: Microbial ecology, v. 47, no. 1, p. 80–
 410 86, doi: 10.1007/s00248-003-1021-z.

411 Wu, H., Chen, W., Rong, X., Cai, P., Dai, K., and Huang, Q., 2014, Soil colloids and
 412 minerals modulate metabolic activity of *Pseudomonas putida* measured using
 413 microcalorimetry: Geomicrobiology Journal, v. 31, p. 590–596, doi:
 414 10.1080/01490451.2013.861544.

Yuan, X., Chen, Z., Xiao, S.H., Zhou, C., Hua, H., 2011, An early Ediacaran assemblage of macroscopic and morphologically differentiated eukaryotes: Nature, v. 470, p. 390-393, doi:10.1038/nature09810.

FIGURE CAPTIONS

Figure 1. Normalized optical density at 600 nm (ΔOD_{600}) of six-hour-old *Pseudoalteromonas luteoviolacea* subcultures taken after 24 hours growth with A: 5, B: 10 and C: 25 mg/ml <2-3 μ m mineral particles (B = berthierine, C = calcite, G = glauconite, I = illite, K = kaolinite) and with no mineral particles (Control). Floating points are outliers. Experiments were conducted in quadruplicate. Values shown represent increases above t=0 and are normalized to the control.





Figure 2 Normalized optical density at 600 nm (ΔOD_{600}) of six-hour-old *Pseudoalteromonas luteoviolacea* subcultures taken after 24 hours growth with 5, 10 and 25 mg/ml <2-3 μ m mineral particles and with no mineral particles (Control). Berthierine, glauconite, illite and kaolinite are grouped together as Clay. Experiments were conducted

432 in quadruplicate. Values shown represent increases above t=0 and are normalised to the
433 control.



434